arXiv:0811.0126v1 [nlin.CD] 2 Nov 2008

On The Geometrical Description of Dynamical Stability

Eduardo Cuervo-Reyes[1,*] and Ramis Movassagh[1,2,†]
[1]*LAC ETH-Zurich, Switzerland*
[2]*Collegium Helveticum, Zurich, Switzerland*
(Dated: November 4, 2008)

A general parametrization requirement for geometrization of dynamical stability is presented. We show that the non-physical behaviors appear in lower dimensional systems when non-affine parametrization of arc length with time is used. We compare the two widely used Jacobi and Eisenhart metrics as archetypes for (non)affine parametrization. We numerically investigate this in the context of the two-centered Morse potential. The relevance of parametric resonance as a source of instability in two dimensional systems is resolved.

PACS numbers:

The general time-independent Hamiltonian

$$H(\boldsymbol{q},\boldsymbol{p})=\frac{1}{2}a^{ij}(\boldsymbol{q})p_ip_j+V(\boldsymbol{q}) \tag{1}$$
$$p_i=a_{ik}(\boldsymbol{q})\dot{q}^k$$

is the basic representation of many problems and applications in the physical sciences ranging from molecular physics to astrophysics. Naturally different approaches for fundamental understanding of such systems and their dynamical (in) stabilities have been the subject of vigorous research [1, 2]. Due to its non-perturbative nature, geometrization of dynamics has been proposed as a novel approach [3, 4], which gives more physically meaningful results for the stability when the traditional Lyapunov exponent does not.[5]

The motion of a mechanical system is entirely determined by the principle of least action [6]. Geometrization of dynamics represents the motion as a geodesic on a manifold with a suitable metric $g=g_{\mu\nu}dq^\mu\otimes dq^\nu$. For a congruence of geodesics $\gamma^i(\tau,s)$, the vector field of geodesic spread is defined as

$$\xi^i(s)=\left[\frac{\partial\gamma^i(\tau,s)}{\partial\tau}\right]_{\tau=\tau_0} \tag{2}$$

where s is the arc length and τ parameterizes the different geodesics. $\xi(s)$ is a measure of the separation of nearby geodesics and its evolution in time or arc-length has been considered as an indicator of (in)stability. This approach can be utilized to tackle a broad range of applications in non-integrable systems and can give new insights to the on-set of chaos. ξ^i evolves according to Jacobi-Levi-Civita (JLC) equation[7], which in local coordinates reads as

$$\frac{D^2\xi^i}{ds^2}+R^i_{jkm}\frac{dq^j}{ds}\xi^k\frac{dq^m}{ds}=0 \tag{3}$$

where $\frac{D}{ds}$ and R^i_{jkm}stand respectively for the covariant derivative along geodesics and the Riemann-Christoffel curvature tensor. Among other possible choices, two metrics that have been extensively used are: the Jacobi or the kinetic energy metric(g_J) on the *admissible* configuration space $M_J=\{x\in M: V(x)<E\}$

$$(g_J)_{ij}\equiv 2[E-V(\boldsymbol{q})]a_{ij}(\boldsymbol{q}) \tag{4}$$

and the Eisenhart metric[8]. The latter is defined on the enlarged ($M\times\Re^2$, g_E) configuration space-time $x^\nu=\{q^0\equiv t,q^i,q^{N+1}\}$ with the arc-length

$$ds^2=-2V(\boldsymbol{q})(dq^0)^2+a_{ij}dq^idq^j+2dq^0dq^{N+1} \tag{5}$$

In the Eisenhart's metric physical motions satisfy an *affine parametrization* $ds^2=\kappa^2dt^2$ (κ is a real constant). The time evolution of the extra coordinate can formally be integrated

$$q^{N+1}=\frac{\kappa^2}{2}t+C_0-\int_0^t\mathcal{L}dt' \tag{6}$$

Equations of motion for ξ using Eisenhart metric are equivalent to those of the tangent dynamics, thus showing its geometrical origin[9]. Solutions of JLC equations are quite complicated and computationally expensive to obtain; they require the simultaneous computation of $q^i(s)$ and $\xi^i(s)$ [10]. For this reason numerical integration of the exact equations (3) has been mostly restricted to low dimensional systems (i.e. two degrees of freedom)[9, 11] and the equivalence of results using both metrics has been argued by simulations of particular systems[9, 12]. In the case of many degrees of freedom, an approximate form of JLC equation for $||\xi||$ has been obtained[10]; this scalar function supposedly gives sufficient information about stability. Moreover, in the thermodynamic limit, under statistical considerations a version of this treatment has been proposed whose main goal is to quantify the stability based only on static geometrical properties [13, 14]. This by-passes the need of computing trajectories for a representative set of initial conditions and gives rise to a potentially powerful tool for finding order-to-chaos transitions. The main argument in favor of using Eisenhart metric up to now is its computational simplicity. On the other hand in the framework of Jacobi metric, the scalar, Ricci's and sectional curvatures have been shown to be positive in many systems of interest [10, 11]. This led to the conclusion that negative curvature is not the fundamental source of instability and to the conjecture that the oscillating curvature leads the systems to chaos through *parametric resonance*. Moreover, Jacobi metric has been considered to be a preferred choice due to its mathematical properties, namely its compactness and Riemannian character[12]. It has been used for stability analysis of a system of self-gravitating particles[10]. Interestingly it was observed that, for the same energy per degree of freedom, increasing the number of particles leads to less divergence in the geodesic spread ξ, namely suppresses chaos. These result of simulations were assumed to be due to the mathematical approximations [10, 11] that were made for obtaining the dynamics of $||\xi||$. Instead, one expects the probability of chaotic dynamics to increase with the number of non-separable degrees of freedom N.

In the present letter we wish to show the fundamentally different results that can emerge from the application of the different metrics mentioned above. We show that in the low dimensional systems non-affine parametrization naturally leads to unstable solutions that should neither be confused with parametric resonance nor be interpreted as physical instabilities. We will discuss conditions under which stability analysis can be performed using either of the two metrics.

A manifold has a well-defined tangent space everywhere. Boundaries of accessible configuration space ∂M_J in Jacobi metric correspond to points where the kinetic energy, T, is zero (i.e. $E = V(q)$). The physical picture is that the geodesics reflect from ∂M_J. Moreover, the reflections have to be perpendicular to ∂M_J otherwise there would be a net momentum parallel to ∂M_J, which would contradict the definition of ∂M_J ($T = 0$)[15, 16]. The curvature tensor in terms of Christoffel symbols is:

$$R^i_{jlk} = \frac{\partial \Gamma^i_{jk}}{\partial q^l} - \frac{\partial \Gamma^i_{jl}}{\partial q^k} + \Gamma^m_{jk}\Gamma^i_{ml} - \Gamma^m_{jl}\Gamma^i_{mk} \tag{7}$$

$$\Gamma^i_{jk} = \frac{1}{2} g^{im}\{g_{jm,k} + g_{km,j} - g_{jk,m}\} \tag{8}$$

Here, g_{ij} is given by Eq.(4); thus Γ^i_{jk} contain powers of $[E - V(q)]^{-1}$ that become infinite on ∂M_J. Clearly this corresponds to cusps in the curvature, where the tangent space is not defined. Moreover there are stationary geodesics at the boundaries, which in general are not equilibrium points. Therefore, these trajectories are not physically meaningful on ∂M_J, in addition to the usage of manifolds not being allowed. However, in the Eisenhart's metric all curvatures are finite (Ricci, sectional, and scalar). From a physical point of view the stability is determined by comparing the difference of trajectories in time, say from the value of Lyapunov exponent.

It is evident from (Eq.2) that ξ by definition compares the geodesics at the same *arc length*. When parametrization is affine, as in Eisenhart's metric, comparing geodesics at equal arc length corresponds to comparison of trajectories at the same time. Therefore, the stability analysis is in a one-to-one correspondence between the divergence of trajectories in time and the geodesic spread.

A natural question is where and under what conditions does the Jacobi metric give similar results for stability analysis to those given by the Eisenhart metric? A first criterion would be to avoid singularities in the curvatures. A conserved nonzero angular momentum for bounded systems warranties that all momentums are *not* simultaneously zero, which puts a lower bound on the kinetic energy. Another favorable condition could be a high number of degrees of freedom, as this generally reduces the oscillations of the kinetic energy. By the arguments given above, the usage of Jacobi metric method for systems with few degrees of freedom and zero total angular momentum should therefore be avoided. A detailed analysis with respect to the dimensionality and coherence of the system will be addressed in a sequel. Here we focus on a two-dimensional problem.



FIG. 1: Poincare maps for $\Delta E = 7.5$

In order to make our analysis more illustrative we apply this formalism to a two-center Morse potential with center-to-center distance $r_e = 3.0271\mathring{A}$, and Morse parameters $D = 40.75$ cm^{-1}, $\alpha = 1.56\mathring{A}^{-1}$, and $\bar{d} = 4.36\mathring{A}$ [9]. The corresponding Hamiltonian with zero total angular momentum is

$$H = \frac{P_R^2}{2\mu_1} + \left(\frac{1}{2\mu_1 R^2} + \frac{1}{2\mu_2 r_e^2}\right) P_\theta^2 + V(R,\theta) \tag{9}$$

where, R and θ are polar coordinates; the polar axis runs across the two centers, and the reduced masses are taken as $\mu_1 = 18$ amu, $\mu_2 = 64$ amu[9]. In the figures, we have expressed all relevant quantities in units that are powers of cm (i.e. energy, linear momentum and λ are in cm^{-1}, t and distances are in cm, thus angular momentum becomes dimensionless). In Fig.1 we show a Poincare Surface Section (PSS) for $\Delta E = E - 2D = 7.5$ cm^{-1} where it is predominated by stable limit-cycles. substituting for $s \rightarrow s(t)$ We plot the Lyapunov exponent $\lambda = \lim_{t\rightarrow\infty} \frac{1}{t}\log(\frac{||\xi(t)||}{||\xi(0)||})$ versus the energy using the Eisenhart metric (Fig.2). λ is a function of energy and the initial conditions; we choose a set of initial conditions in the semi-stable subspace for $\Delta E = 7.5$ cm^{-1} in agreement with Fig.1. As one expects the lower energy values correspond to stable solutions whereas the higher energies, $\Delta E > 4.3$ cm^{-1}, produce positive values of λ and therefore, the solutions exponentially diverge. In order to ensure a proper convergence to the Lyapunov exponent we show in Fig.3 the evolution of $Y(t) = \frac{1}{t}\log(\frac{||\xi(t)||}{||\xi(0)||})$ for different energies ranging from very stable ($\Delta E = 0.5$ cm^{-1}) to unstable ($\Delta E = 15.5$ cm^{-1}) regimes. It is important to note that the stable dynamics demands $\lambda \leq 0$. The same is shown in Fig.4 for a very stable regime ($\Delta E = 2.5$ cm^{-1}) using Jacobi metric. As can be seen, Y converges to about 130 cm^{-1}, in contradiction to the PSS and to the results obtained from Eisenhart metric (see Fig.2)! We also plotted in Fig.5 the behavior of the Y function in a small time window for the same initial conditions and energy as in Fig.4. It is clear that the nonphysical limit for the convergence of the Y-function in Fig.4 is a consequence of the "kicks" produced by the oscillations in the kinetic energy. Although we have chosen as an example a particular system, the appearance of singularities in the curvature and its oscillations are general consequences of the Jacobi metric. In this case, the curvature is actually positive and therefore the exponential growth of ξ is due to the dominant quasi-periodic "kicks" produced by the kinetic energy values close to zero as is evident in (equations 3, and 7).



FIG. 2: Lyapunov values versus Energy using Eisenhart's metric



FIG. 3: Lyapunov convergence for different energies using Eisenhart metric



FIG. 4: Kinetic Energy and Lyapunov convergence for ΔE=2.5 cm^{-1} using Jacobi metric

Before concluding we make some final remarks to put our work into a larger context of recent developments. Though still mathematically unproven we have shown a counter-example against the reliability of the Jacobi metric for geometrization of dynamical stability. The reason this has gone unnoticed in the past is most probably due to the fact that the formalisms were applied in regimes, where i) Jacobi metric could be approximated as affine, i.e. small changes of the kinetic energy, as can be the case for large number of degrees of freedom[12, 13, 17, 18] ii) the particular conditions, under which comparison between tangent dynamics and the Jacobi metric were done, were already in the chaotic regime or at the onset of chaos; thus the discrepancy between Eisenhart and Jacobi metrics could not be clearly established[10, 11] iii) calculations were done mostly using the Eisenhart met-



FIG. 5: Zooming in on Fig.4: Kinetic Energy and the corresponding "kicks" in ξ for ΔE=2.5 cm^{-1}

ric and assumed to hold for Jacobi as well (please see references in [12]). Furthermore, parametric resonance is a very particular mechanism for driving a system to instability (please see Landau and Lifshitz [6]), which in the past has not been firmly investigated as a source of chaos within the Jacobi framework. Here we show how semi-periodic kicks produced by small values of T, which is a side-effect of the usage of the Jacobi metric as opposed to intrinsic dynamical property of the system, may be misinterpreted as actual physical parametric resonance. To our knowledge Eisenhart's metric remains to be a very successful method for geometrization of general holonomic conservative systems in classical dynamics. In addition geometrizatoin is much easier to work with using Eisenhart's metric.

In this letter we have discussed an important requirement for geometrization of stability analysis. We have explicitly shown that the time-evolution of the geodesic spread, in lower dimensional systems, is not a measure of stability when the parametrization of the arc length with time is non-affine. Due to their importance, we put this discussion in the context of Jacobi and Eisenhart metrics, which can be looked at as the archetypes of non- and affine parametrizations respectively. The apparent parametric resonance phenomena seen in the frame-work of Jacobi geometrization discussed in earlier works is most probably due to the oscillations of the kinetic energy regardless of the stability of the system. Future work entails a thorough investigation of stability for higher degrees of freedom systems as well as the explicit dependence of the stability on the degree of non-affinity-the fluctuations in the kinetic energy.

We thank Reinhard Nesper, Richard Rand, Reinel Sospedra-Alfonso for discussions. This project was funded by Swiss National Science Foundation and The Cogito Foundation.

---

[*] E. Cuervo-Reyes and Ramis Movassagh contributed equally to this work

[†] Corresponding author: movassagh@collegium.ethz.ch

[1] H. Poincaré, *Les Méthodes Nouvelles de la Méchanique Celeste* Blanchard, Paris, vol. 3 (1987).

[2] V. I. Arnold, *Mathematical methods for classical mechanics* (MIR, Moscow, 1976).

[3] N. S. Krylov, *Works on the Fundations of Statistical Mechanics* (Princeton Univ. Press, Princeton, 1979)

[4] M. Pettini, Phys. Rev. E **47**, 828 (1993).

[5] R. Rand, Int. J. Non-linear Mechanics **8** pp.161-168,(1973)

[6] L. D. Landau and E. M. Lifshitz *Mechanics* (1976).

[7] T. Levi-Civita, Ann. Math. 97, 291, (1926).

[8] L. P. Eisenhart, Ann. Math. **30**, 591 (1929)

[9] R. Sospedra-Alfonso, L. Velazquez, and J. Rubayo-Soneira Chem. Phys. Lett. 37 261-268 (2003).

[10] M. Cerruti-Sola and M. Pettini, Phys. Rev. E **51**, 53-64 (1995).

[11] M. Cerruti-Sola and M. Pettini Phys. Rev. E **53**, 179-188 (1996)

[12] M. Cerruti-Sola, R. Franzosi and M. Pettini, Phys. Rev. E **56**, 4872-4875 (1997).

[13] L. Casetti, R. Livi and M. Pettini Phys. Rev. Lett. **74**, 375-378 (1995)

[14] L. Caiani, L. Casetti, C. Clementi and M. Pettini Phys. Rev. Lett. **79**, 4361-4364 (1997).

[15] M. Szydlowski, M. Heller, W. Sasin, J. Math. Phys. **37**, 1 345 January (1996).

[16] M. Szydlowski, J. Math. Phys. **40**, 7 345 July (1999).

[17] L. Casetti, C. Clementi and M. Pettini Phys. Rev. E **54**, 5969-5984 (1996).

[18] L. Casetti, M. Pettini, E.G.D. Cohen Phys. Rep. **337**, 237-341 (2000).